

Vaccine Supplies

Vaccine	Route/Needle	Notes	Supplies
Hepatitis A and B Brand: Twinrix®	IM- See Below	<ul style="list-style-type: none"> IM injection Standard dosing: series of three doses (0-, 1-, and 6-months) 1mL doses Tip caps of prefilled syringes contain rubber latex Hazy white suspension Shake the syringe by tipping it upside down and then back upright for at least 15 seconds before administering 	<ul style="list-style-type: none"> Standard supplies* + Needle
Hepatitis B (HBV) Brand: Engerix-B®	IM- See Below	<ul style="list-style-type: none"> IM injection Standard dosing: series of three doses (0-, 1-, and 6-months) 19-years and younger receive 0.5mL doses 20-years and older receive 1mL doses Tip caps of prefilled syringes contain rubber latex Turbid white suspension Shake well before administering 	<ul style="list-style-type: none"> Standard supplies* + Needle
Influenza Brands: Fluarix®, Flucelvax®, Flulaval®, Fluzone HD®	IM- See Below	<ul style="list-style-type: none"> Please refer to yearly influenza vaccine information handout. Note: Flucelvax® contains NO egg protein or antibiotics 	<ul style="list-style-type: none"> Standard supplies* + Needle
Pneumococcal PPSV23 Brand: Pneumovax® 23	IM- See Below	<ul style="list-style-type: none"> IM or Subcutaneous techniques are appropriate Single 0.5 mL dose If given subcutaneously, use a 5/8", 23-25-gauge needle Clear, sterile solution 	<ul style="list-style-type: none"> Standard supplies* + Needle
Pneumococcal PCV13 Brand: Prevnar 13®	IM- See Below	<ul style="list-style-type: none"> IM only Single 0.5mL dose Tip caps of prefilled syringes contain rubber latex White suspension Shake well immediately before use 	<ul style="list-style-type: none"> Standard supplies* + Needle
Tdap Brand: Boostrix®	IM- See Below	<ul style="list-style-type: none"> IM injection Single 0.5mL dose Tip caps of prefilled syringes contain rubber latex Turbid white suspension Shake well before administering 	<ul style="list-style-type: none"> Standard supplies* + Needle

Vaccine Supplies

Vaccine	Route/Needle	Notes	Supplies
Zoster Brand: SHINGRIX	IM- See Below	<ul style="list-style-type: none"> IM injection Standard dosing: series of two doses (0- and 2-6 months) Reconstituted doses are 0.5mL Supplied in two vials to be combined before administration Sterile white powder (antigen component) and opalescent, colorless to pale brown liquid (adjuvant suspension) 	<ul style="list-style-type: none"> Standard supplies* + TWO needles (one for reconstitution and one for administration) One syringe with at least 0.5mL volume capacity Reconstituted; Give BOTH the powder and liquid vials to provider
Tuberculin Brand: Tubersol®	<ul style="list-style-type: none"> 27-gauge with short bevel ¼- ½ Inch 	<ul style="list-style-type: none"> Intradermal placement Protect from light A vial entered and in use for 30 days should be discarded Clear, colorless liquid 	<ul style="list-style-type: none"> Standard supplies* + Syringe and needle labeled as "TB Syringe"

Intramuscular (IM) Injections			
Gender and Weight	Needle Gauge	Needle Length (inches)	Injection Site
Female or male less than 130 lbs.	22-25	5/8"-1"	Deltoid muscle of arm
Female or male 130-152 lbs.	22-25	1"	Deltoid muscle of arm
Female 153-200 lbs.	22-25	1-1½"	Deltoid muscle of arm
Male 153-260 lbs.	22-25	1-1½"	Deltoid muscle of arm
Female 200+ lbs.	22-25	1½"	Deltoid muscle of arm
Male 260+ lbs.	22-25	1½"	Deltoid muscle of arm

*Standard supplies: paper towels, alcohol prep pads, cotton gauze, band aids, gloves, sharps container

Please ensure providers have all necessary supplies. Please note, except for SHINGRIX and TB skin tests, supply requirements assume prefilled syringes are available for vaccine doses. If not, please issue an additional 1mL syringe and one additional needle.